

Corollary 4. $P_{\mathbf{x}}^*(\mathbf{y})$ is transportable from Π to Π^* in G if and only if there is not s -hedge for $P_{\mathbf{x}'}^*(\mathbf{y}')$ in G for any $\mathbf{X}' \subseteq \mathbf{X}$ and $\mathbf{Y}' \subseteq \mathbf{Y}$.

Proof. See Appendix 2. ■

Theorem 7. The rules of do-calculus, together with standard probability manipulations are complete for establishing transportability of all effects of the form $P_{\mathbf{x}}^*(\mathbf{y})$.

Proof. See Appendix 2. ■

6 Other perspectives on generalizability

Many problems in statistics and causal inference can be framed as problems of generalizability, though inherently different from that of transportability.

Consider, for example, classical statistical inference, it can be viewed as a generalization from properties of a *random* sample Π_S of a population Π to properties of the population Π itself. Two centuries of statistical analysis have rendered this task well understood and fairly complete.

Next consider the problem of causal inference, that is, to estimate causal-effects from observational studies (given a set of causal assumptions). This class of problems can be viewed as a generalization from a population under observational regime to a population under experimental regime. Since the imposition of experimental regime (e.g., forcing individuals to receive treatment) induces a behavioral change in the population, the problem can be viewed as generalization between two diverse populations. Fortunately, the disparities between the two populations are local (assumes atomic interventions), involving only the treatment assignment mechanism and, so, with the help of model assumptions, a complete solution to the problem can be obtained (using do-calculus). We can decide algorithmically whether the assumptions at hand are sufficient for estimating a given causal effect and, if the answer is affirmative, we can derive its estimand.

An important variant in causal inference is the task of estimating causal effects from surrogate experiments, namely, experiments in which a surrogate set of variables Z are manipulated, rather than the one (X) whose effect we seek to estimate.¹¹ This variant too can be viewed as an exercise in generalization, this time from a population under regime $do(Z = z)$ to that same population under regime $do(X = x)$. A complete solution to this problem is reported in [37].

Another challenge of generalizability flavor arises, in both observational and experimental studies, when samples Π_S are not randomly drawn from the population of interest Π , but are selected preferentially, depending on the values taken by a set V_S of variables. This problem, known as “selection bias” (or “sampling selection bias”), has received due attention in epidemiology, statistics, and economics [38–41] and can be viewed as a generalization from the sampled population to the population at large, when little is known about their relationships save for qualitative assumptions about the selection mechanism. Graphical models were used to improve the understanding of the problem [42–45] and gave rise to several conditions for recovering from selection bias when the probability of selection is available.

Likewise, Refs. 21, 46, 47 tackle variants of the sample selection problem assuming that certain relationships are invariant between the two groups (i.e., sample and population). The former assumed knowledge of the probability of selection in each of the principal stratum, while the latter exploited (using propensity score analysis) the availability of the probability of selection in each combination of covariates.

¹¹ A surrogate variable is different from instrumental variable in that the former should lead to the identification of causal effect even in nonparametric models; IV methods are limited to “local” causal effects (so-called LATE [48]).

More recently, Didelez et al. [49] studied conditions for recovering from selection bias when no quantitative knowledge is available about selection probabilities. Bareinboim and Pearl [50] extended these conditions and provided a complete characterization, together with an algorithm, for deciding when a bias-free estimate of the odds ratio (OR) can be recovered from selection-biased data. They also developed methods using instrumental variables that recover other effect measures when information about the target population is available for some variables (see also Ref. 51).

The problem of transportability is fundamentally different from the other problems of generalizability discussed above. Transportability deals with two distinct populations that are different both in their inherent characteristics (encoded by the S variables) and the regimes under which they are studied (i.e., experimental vs. observational).

Hernán and VanderWeele [52] addressed a problem related to transportability in the context of “compound treatments,” namely, treatments that can be implemented in multiple versions (e.g., “exercise at least 15 minutes a day”). Transportability arises when we wish to predict the response of a population that implements one version of the treatment from a study on another population, in which another version is implemented. Petersen [53] showed that this problem is a variant of the general problem treated in Ref. 1, to which this article provides an algorithmic solution.

Finally, it is important to mention two recent extensions of the results reported in this article. Bareinboim and Pearl [2] have addressed the problem of transportability in cases where only a limited set of experiments can be conducted at the source environment. Subsequently, the results were generalized to the problem of “meta-transportability,” that is, pooling experimental results from multiple and disparate sources to synthesize a consistent estimate of a causal relation at yet another environment, potentially different from each of the formers [3].

7 Conclusions

Informal discussions concerning the difficulties of generalizing experimental results across populations have been going on for almost half a century [4, 5, 54–56] and appear to accompany every textbook in experimental design. By and large, these discussions have led to the obvious conclusions that researchers should be extremely cautious about unwarranted generalization, that many threats may await the unwary, and that extrapolation across studies requires “some understanding of the reasons for the differences” [54, p. 11].

The formalization offered in this article embeds this discussion in a precise mathematical language and provides researchers with theoretical guarantees that, if certain conditions can be ascertained, generalization across populations can be accomplished, protected from the threats and dangers that the informal literature has accumulated.

Given judgmental assessments of how target populations may differ from those under study, the article offers a formal representational language for making these assessments precise (Definition 3) and, subsequently, deciding whether, and how, causal relations in the target population can be inferred from those obtained in experimental studies. Corollary 4 in this article provides a complete (necessary and sufficient) graphical condition for deciding this question and, whenever satisfied, we further provide an algorithm for computing the correct transport formula (Figure 5). The transport formula specifies the proper way of modifying the experimental results so as to account for differences in the populations. These transport formulae enable the investigator to select the essential measurements in both the experimental and observational studies and combine them into a bias-free estimand of the target quantity.

While the results of this article concern the transfer of causal information from experimental to observational studies, the method can also benefit in transporting statistical findings from one observational study to another [57]. The rationale for such transfer is twofold. First, information from the first study may enable researchers to avoid repeated measurement of certain variables in the target population.

Second, by pooling data from both populations, we increase the precision in which their commonalities are estimated and, indirectly, also increase the precision by which the target relationship is transported. Substantial reduction in sampling variability can be thus achieved through this decomposition [58].

Of course, our analysis is based on the assumption that the analyst is in possession of sufficient background knowledge to determine, at least qualitatively, *where* two populations may differ from one another. In practice, such knowledge may only be partially available. Still, as in every mathematical exercise, the benefit of the analysis lies primarily in understanding what must be assumed about reality for generalization to be valid, what knowledge is needed for a given task to succeed, and how sensitive conclusions are to knowledge that we do not possess.

Acknowledgment: A preliminary version of this article was presented at the 26th AAAI Conference, Toronto, CA, July, 2012 [59]. We appreciate the insightful comments provided by two anonymous referees. This article benefited from discussions with Onyebuchi Arah, Stuart Baker, Susan Ellenberg, Eleazar Eskin, Constantine Frangakis, Sander Greenland, David Heckerman, James Heckman, Michael Hoefler, Marshall Joffe, Rosa Matzkin, Geert Molengergh, William Shadish, Ian Shrier, Dylan Small, Corwin Zigler, and Song-Chun Zhu.

This research was supported in parts by grants from NSF #IIS-1249822, and ONR #N00014-13-1-0153 and #N00014-10-1-0933.

Appendix 1: causal assumptions in nonparametric models

The tools presented in this article were developed in the framework of nonparametric SCM, which subsumes and unifies many approaches to causal inference.¹²

A SCM M conveys a set of assumptions about how the world operates. This contrasts the statistical tradition in which a model is defined as a set of distributions (see footnote 15). Causal models is better viewed as a set of assumptions about Nature, with the understanding that each assumption (i.e., that the set of arguments of f_i does not include variable V_j) constrains the set of distributions (like $P(v)$) that the model can generate.

The formal structure of SCM's was defined in Section 3, here we illustrate their power as inference engines. Consider a simple SCM model depicted in Figure 6(a), which represents the following three functions:

$$\begin{aligned} z &= f_Z(u_Z) \\ x &= f_X(z, u_X) \\ y &= f_Y(x, u_Y), \end{aligned} \tag{10}$$

where in this particular example, U_Z , U_X , and U_Y are assumed to be jointly independent but otherwise arbitrarily distributed. Each of these functions represents a causal process (or mechanism) that determines




Figure 6 The diagrams associated with (a) the structural model of eq. [6] and (b) the modified model of eq. [11], representing the intervention $do(X = x_0)$.

¹² We use the acronym SCM for both parametric and non-parametric representations (which is also called Structural Equation Model (SEM)), though historically, SEM practitioners preferred the parametric representation and often confuse with regression equations [60].

the value of the left variable (output) from the values on the right variables (inputs) and is assumed to be invariant unless explicitly intervened on. The absence of a variable from the right-hand side of an equation encodes the assumption that nature ignores that variable in the process of determining the value of the output variable. For example, the absence of variable Z from the arguments of f_Y conveys the empirical claim that variations in Z will leave Y unchanged, as long as variables U_Y and X remain constant.

Representing Interventions, counterfactuals, and causal effects

This feature of invariance permits us to derive powerful claims about causal effects and counterfactuals, even in nonparametric models, where all functions and distributions remain unknown. This is done through a mathematical operator called $do(x)$, which simulates physical interventions by deleting certain functions from the model, replacing them with a constant $X = x$, while keeping the rest of the model unchanged [61–63]. For example, to emulate an intervention $do(x_0)$ that holds X constant (at $X = x_0$) in model M of Figure 6(a), we replace the equation for x in eq. [10] with $x = x_0$, and obtain a new model, M_{x_0} ,

$$\begin{aligned} z &= f_Z(u_Z) \\ x &= x_0 \\ y &= f_Y(x, u_Y), \end{aligned} \quad [11]$$

the graphical description of which is shown in Figure 6(b).

The joint distribution associated with the modified model, denoted $P(z, y|do(x_0))$ describes the post-intervention distribution of variables Y and Z (also called “controlled” or “experimental” distribution), to be distinguished from the preintervention distribution, $P(x, y, z)$, associated with the original model of eq. [10]. For example, if X represents a treatment variable, Y a response variable, and Z some covariate that affects the amount of treatment received, then the distribution $P(z, y|do(x_0))$ gives the proportion of individuals that would attain response level $Y = y$ and covariate level $Z = z$ under the hypothetical situation in which treatment $X = x_0$ is administered uniformly to the population.¹³

In general, we can formally define the postintervention distribution by the equation

$$P_M(y|do(x)) = P_{M_x}(y) \quad [12]$$

In words, in the framework of model M , the postintervention distribution of outcome Y is defined as the probability that model M_x assigns to each outcome level $Y = y$. From this distribution, which is readily computed from any fully specified model M , we are able to assess treatment efficacy by comparing aspects of this distribution at different levels of x_0 .¹⁴

Identification, d-separation and causal calculus

A central question in causal analysis is the question of *identification* in partially specified models: Given assumptions set A (as embodied in the model), can the controlled (postintervention) distribution, $P(y|do(x))$, be estimated from data governed by the preintervention distribution $P(z, x, y)$?

In linear parametric settings, the question of identification reduces to asking whether some model parameter, β , has a unique solution in terms of the parameters of P (say the population covariance matrix).

¹³ Equivalently, $P(z, y|do(x_0))$ can be interpreted as the joint probability of $(Z = z, Y = y)$ under a randomized experiment among units receiving treatment level $X = x_0$. Readers versed in potential-outcome notations may interpret $P(y|do(x), z)$ as the probability $P(Y_x = y|Z_x = z)$, where Y_x is the potential outcome under treatment $X = x$.

¹⁴ Counterfactuals are defined similarly through the equation $Y_x(u) = Y_{M_x}(u)$ (see [16, Ch. 7]), but will not be needed for the discussions in this article.

In the nonparametric formulation, the notion of “has a unique solution” does not directly apply since quantities such as $Q(M) = P(y|do(x))$ have no parametric signature and are defined procedurally by simulating an intervention in a causal model M , as in eq. [11]. The following definition captures the requirement that Q be estimable from the data:

Definition 11 (Identifiability).¹⁵ *A causal query $Q(M)$ is identifiable, given a set of assumptions A , if for any two models (fully specified) M_1 and M_2 that satisfy A , we have*

$$P(M_1) = P(M_2) \Rightarrow Q(M_1) = Q(M_2) \quad [13]$$

In words, the functional details of M_1 and M_2 do not matter; what matters is that the assumptions in A (e.g., those encoded in the diagram) would constrain the variability of those details in such a way that equality of P 's would entail equality of Q 's. When this happens, Q depends on P only, and should therefore be expressible in terms of the parameters of P .

When a query Q is given in the form of a do-expression, for example $Q = P(y|do(x), z)$, its identifiability can be decided systematically using an algebraic procedure known as the do-calculus [13]. It consists of three inference rules that permit us to map interventional and observational distributions whenever certain conditions hold in the causal diagram G .

The conditions that permit the application these inference rules can be read off the diagrams using a graphical criterion known as d -separation [65].

Definition 12 (d -separation). *A set S of nodes is said to block a path p if either*

1. p contains at least one arrow-emitting node that is in S , or
2. p contains at least one collision node that is outside S and has no descendant in S .

If S blocks all paths from set X to set Y , it is said to “ d -separate X and Y ,” and then, it can be shown that variables X and Y are independent given S , written $X \perp\!\!\!\perp Y|S$.¹⁶

D -separation reflects conditional independencies that hold in any distribution $P(v)$ that is compatible with the causal assumptions A embedded in the diagram. To illustrate, the path $U_Z \rightarrow Z \rightarrow X \rightarrow Y$ in Figure 6(a) is blocked by $S = \{Z\}$ and by $S = \{X\}$, since each emits an arrow along that path. Consequently we can infer that the conditional independencies $U_Z \perp\!\!\!\perp Y|Z$ and $U_Z \perp\!\!\!\perp Y|X$ will be satisfied in any probability function that this model can generate, regardless of how we parametrize the arrows. Likewise, the path $U_Z \rightarrow Z \rightarrow X \leftarrow U_X$ is blocked by the null set $\{\emptyset\}$, but it is not blocked by $S = \{Y\}$ since Y is a descendant of the collision node X . Consequently, the marginal independence $U_Z \perp\!\!\!\perp U_X$ will hold in the distribution, but $U_Z \perp\!\!\!\perp U_X|Y$ may or may not hold.¹⁷

The rules of *do*-calculus

Let X , Y , Z , and W be arbitrary disjoint sets of nodes in a causal DAG G . We denote by $G_{\overline{X}}$ the graph obtained by deleting from G all arrows pointing to nodes in X . Likewise, we denote by $G_{\underline{X}}$ the graph obtained by

¹⁵ This definition appears to be similar to, but differ fundamentally from the standard statistical definition [64, p. 22] which deals with the unidentifiability of the parameter set θ from a distribution P_θ . In our case, the query $Q = P(Y|do(x))$ is not a parameter of P (see [22, p. 77]).

¹⁶ See Hayduk et al. [66], Glymour and Greenland [67], and Pearl [16, p. 335] for a gentle introduction to d -separation.

¹⁷ This special handling of collision nodes (or *colliders*, e.g., $Z \rightarrow X \leftarrow U_X$) reflects a general phenomenon known as *Berkson's paradox* [68], whereby observations on a common consequence of two independent causes render those causes dependent. For example, the outcomes of two independent coins are rendered dependent by the testimony that at least one of them is a tail.

deleting from G all arrows emerging from nodes in X . To represent the deletion of both incoming and outgoing arrows, we use the notation $G_{\overline{XZ}}$.

The following three rules are valid for every interventional distribution compatible with G .

Rule 1 (Insertion/deletion of observations):

$$P(y|do(x), z, w) = P(y|do(x), w) \text{ if } (Y \perp\!\!\!\perp Z|X, W)_{G_{\overline{X}}} \quad [14]$$

Rule 2 (Action/observation exchange):

$$P(y|do(x), do(z), w) = P(y|do(x), z, w) \text{ if } (Y \perp\!\!\!\perp Z|X, W)_{G_{\overline{XZ}}} \quad [15]$$

Rule 3 (Insertion/deletion of actions):

$$P(y|do(x), do(z), w) = P(y|do(x), w) \text{ if } (Y \perp\!\!\!\perp Z|X, W)_{G_{\overline{XZ(W)}}}, \quad [16]$$

where $Z(W)$ is the set of Z -nodes that are not ancestors of any W -node in $G_{\overline{X}}$.

To establish identifiability of a query Q , one needs to repeatedly apply the rules of do-calculus to Q , until the final expression no longer contains a do-operator¹⁸; this renders it estimable from non-experimental data. The do-calculus was proven to be complete to the identifiability of causal effects in the form $Q = P(y|do(x), z)$ [69, 70], which means that if Q cannot be expressed in terms of the probability of observables P by repeated application of these three rules, such an expression does not exist.

We shall see that, to establish transportability, the goal will be different; instead of eliminating do-operators, we will need to separate them from a set of variables S that represent disparities between populations.

Appendix 2

Theorem 2. *Let G be a selection diagram. Then for any node Y , the direct effect $P_{Pa(Y)}^*(y)$ is transportable if there is no subgraph of G which forms a Y -rooted sC -tree.*

Proof. We know from Tian [71, Theorem 22] that whenever there exists no subgraph G_T of G satisfying all of the following: (i) $Y \in T$; (ii) G_T has only one c -component, T itself; (iii) All variables in T are ancestors of Y in G_T , the direct effect on Y is identifiable, as sC -trees are structures of this type. Further Shpitser and Pearl [27, Theorem 2] showed that the same holds for C -trees, which also implies the inexistence of a sC -trees. Since such structure does not show up in G , the target quantity is identifiable, and hence transportable.

It remains to show that the same holds whenever there exists a subgraph that is a C -tree and in which no S node points to Y , i.e., there is no Y -rooted sC -tree at all. It is true that $(S \perp\!\!\!\perp Y|Pa(Y))_{G_{\overline{Pa(Y)}}}$, given that all directed paths from S to Y are closed. This follows from the following facts: (1) all paths from S passing through Y 's ancestors were cut in $G_{\overline{Pa(Y)}}$; (2) all bidirected paths were also closed given that the conditioning set contains only root nodes, and a connection from S must pass through at least one collider; (3) transportability does not depend on descendants of Y (by argument similar to Tian [71, Lemma 9]). Thus, it follows that we can write $P_{Pa(Y)}^*(Y) = P_{Pa(Y)}(Y|S) = P_{Pa(Y)}(Y)$, concluding the proof. ■

Corollary 1. *Let G be a selection diagram. Then for any node Y , the direct effect $P_{Pa(Y)}^*(y)$ is transportable if there is no S node pointing to Y .*

Proof. Follows directly from Theorem 2. ■

¹⁸ Such derivations are illustrated in graphical details in Ref. [16, p. 87].

Lemma 5. *The exclusive OR (XOR) function is commutative and associative.*

Proof. Follows directly from the definition of the XOR function. ■

Remark 1. The construction given below is a strict generalization of Theorem 1, and it is useful because it will provide a simplified construction of the one provided in Theorem 1, and also set the tone for proofs of generic graph structures which will in the sequel show to be instrumental in proving non-transportability in arbitrary structures.

Theorem 3. *Let G be a Y -rooted sC -tree. Then the effects of any set of nodes in G on Y are not transportable.*

Proof. The proof will proceed by constructing a family of counterexamples. For any such G and any set \mathbf{X} , we will construct two causal models M_1 and M_2 that will agree on $\langle P, P^*, I \rangle$, but disagree on the interventional distribution $P_x^*(y)$.

Let the two models M_1, M_2 agree on the following features. All variables in $\mathbf{U} \cup \mathbf{V}$ are binary. All exogenous variables are distributed uniformly. All endogenous variables except Y are set to the bit parity (sum mod 2) of the values of their parents. The two models differ in respect to Y 's definition. Consider the function for Y , $f_Y : U, Pa(Y) \rightarrow Y$ to be defined as follows:

$$\begin{cases} M_1 : Y = ((pa(Y) \otimes u) \otimes s) \\ M_2 : Y = ((pa(Y) \otimes u) \vee s) \end{cases}$$

Lemma 6. *The two models agree in the distributions $\langle P, P^*, I \rangle$.*

Proof. Since the two models agree on $P(U)$ and all functions except f_Y , it suffices to show that f_Y maintains the same input/output behavior in both models for each domains.

Subclaim 1: Let us show that both models agree in the observational and interventional distributions relative to domain Π , i.e., the pair $\langle P, I \rangle$. The index variable S is set to 0 in Π , and f_Y evaluates to $(pa(Y) \otimes u)$ in both models, which proves the subclaim.

Subclaim 2: Let us show that both models agree in the observational distribution relative to Π^* , i.e., P^* . The index variable S is set 1 in Π^* , and f_Y evaluates to $((pa(Y) \otimes u) \otimes 1)$ in M_1 , and 1 in M_2 . Since the evaluation in M_1 can be rewritten as $\neg((pa(Y) \otimes u))$, it remains to show that $(pa(Y) \otimes u)$ always evaluates to 0.

This fact is certainly true, consider the following observations: a) each variable in U has exactly two endogenous children; b) the given tree has Y as the root; c) all functions are XOR – these imply that Y is computing the bit parity of the sum of all U nodes, which turns out to be even, and so evaluates to 0 and proves the subclaim. ■

Lemma 7. *For any set \mathbf{X} , $P_1(Y|do(\mathbf{X}), S = 1) \neq P_2(Y|do(\mathbf{X}), S = 1)$.*

Proof. Given the functional description and the discussion in the previous Lemma, the function f_Y evaluates always to 1 in M_2 .

Now let us consider M_1 . Note that performing the intervention and cutting the edges going toward \mathbf{X} creates an asymmetry on the sum of the bidirected edges departing from \mathbf{U} , and consequently in the sum performed by Y . It will be the case that some \mathbf{U}' will appear only once in the expression of Y . Therefore, depending on the assignment $\mathbf{X} = \mathbf{x}$, we will need to evaluate the sum (mod 2) over \mathbf{U}' in Y or its negation, which given the uniformity of the distribution of \mathbf{U} will yield $P_1(Y|do(\mathbf{X}), S = 1) = 1/2$ in both cases. ■

By Lemma 2, Lemmas 6 and 7 together prove Theorem 3. ■

Corollary 2. Let G be a selection diagram, let \mathbf{X} and \mathbf{Y} be set of variables. If there exists a node W which is an ancestor of some node $Y \in \mathbf{Y}$ and such that there exists a W -rooted sC-tree which contains any variables in \mathbf{X} , then $P_{\mathbf{x}}^*(\mathbf{y})$ is not transportable.

Proof. Fix a W -rooted sC-tree T , and a path p from W to Y . Consider the graph $p \cup T$. Note that in this graph $P_{\mathbf{x}}^*(Y) = \sum_w P_{\mathbf{x}}^*(w)P^*(Y|w)$. From the last Theorem $P_{\mathbf{x}}^*(w)$ is not transportable, it is now easy to construct $P^*(Y|W)$ in such a way that the mapping from $P_{\mathbf{x}}(W)$ to $P_{\mathbf{x}}(\mathbf{Y})$ is one to one, while making sure all distributions are positive.

Remark 2. The previous results comprised cases in which there exist sC-trees involved in the non-transportability of Y – i.e., Y or some of its ancestors were roots of a given sC-tree. In the problem of identifiability, the counterpart of sC-trees (i.e., C-trees) suffices to characterize non-identifiability for singleton Y . But transportability is more subtle and this is not the case here – it not only depends on X and Y “locations” in the graph, but also the relative position of the S-nodes. Consider Figures 4 and 7(a) (called sp-graph). In these graphs there is no sC-tree but the effect of X on Y is still non-transportable.

The main technical subtlety here is that in sC-trees, a S-node combines its effect with a X-node intersecting in the root node (considering only the bidirected edges), which is not the case for non-transportability in general. Note that in the graphs in Figure 4, and the sp-graph, the nodes S and X intersect first through ordinary edges and meet through bidirected edges only on the Y node. This implies a certain “asynchrony” because, in the structural sense, the existence of a S-node implies a difference in the structural equations between domains, but only this difference does not imply non-transportability (for instance, $P_{\mathbf{x}}^*(z)$ is transportable in the sp-graph even though the equations of Z being different in both models).




Figure 7 Selection diagrams in which $P^*(y|do(x))$ is not transportable, there is no sC-tree but there is a sC-forest. These diagrams will be used as basis for the general case; the first diagram is named sp-graph and the second one sb-graph.

The key idea to produce a proof for non-transportability in these cases is to keep the effect of S-nodes after intersecting with X “dormant” until they reach the target Y and then manifest. We implement this idea in the next two proofs, which can be seen as base cases, and should pavement the way for the most general problem.

Theorem 8. $P_{\mathbf{x}}^*(y)$ is not transportable in the sp-graph (Figure 7(a)).

Proof. We will construct two causal models M_1 and M_2 compatible with the sp-graph that will agree on $\langle P, P^*, I \rangle$, but disagree on the interventional distribution $P_{\mathbf{x}}^*(y)$.

Let us assume that all variables in $U \cup V$ are binary, and let U_1 be the common cause of X and Y , U_2 be the common cause of Z and Y , and U_3 be the random disturbance exclusive to Z . Let M_1 and M_2 be defined as follows:

$$M_1 = \begin{cases} X = U_1 \\ Z = (((X \otimes U_2 \otimes 1) \otimes U_3) \vee S) \otimes (S \wedge (X \otimes U_2)) \\ Y = Z \otimes U_1 \otimes U_2 \end{cases}$$

and:

$$M_2 = \begin{cases} X = U_1 \\ Z = (((U_2 \otimes 1) \otimes U_3) \vee S) \otimes (S \wedge U_2) \\ Y = Z \otimes U_2 \end{cases}$$

Both models agree in respect to $P(\mathbf{U})$, which is defined as follows: $P(U_1) = P(U_2) = P(U_3) = 1/2$.

Lemma 8. *The two models agree in the distributions $\langle P, P^*, I \rangle$.*

Proof. Subclaim 1: Let us show that both models agree in the observational and interventional distributions relative to domain Π , i.e., the pair $\langle P, I \rangle$. In both models X has the same expression, which entails the same (uniform) probabilistic behavior in both cases. The index variable S is set to 0 in Π , and Z evaluates to $(X \otimes U_2 \otimes 1 \otimes U_3)$ in M_1 and $(U_2 \otimes 1 \otimes U_3)$ in M_2 . Clearly, for any value of $X = x$, since U is the same and uniformly distributed in both models, we obtain the same (uniform) input/output probabilistic behavior in M_1 and M_2 (note that U_2, U_3 can freely vary independently of X). In similar way, Y evaluates to $(1 + U_3)$ in both models, which entails the same (uniform) input/output probabilistic behavior in both models. In regard to $do(X = x)$, it is clear that Z did not depend (probabilistically) on the specific value of X , and so the equality between both models follows. For the case when we have $do(Z = z)$, Y evaluates to $(Z \otimes U_1 \otimes U_2)$ in M_1 and $(Z \otimes U_2)$ in M_2 , and given the uniformity of U , they preserve the same (uniform) input/output probabilistic behavior. (For a more elaborated argument, see Theorem 4 below.)

Subclaim 2: Let us show that both models agree in the observational distribution P^* relative to Π^* . The index variable S is set 1 in Π^* , f_Z evaluates to $(X \otimes U_2 \otimes 1)$ in M_1 , and $(U_2 \otimes 1)$ in M_2 . Again, for any value of X , together with the uniformity of U , we obtain the same (uniform) input/output probabilistic behavior in both models (note again that U_2 can freely vary independently of variations of X , and so Z). Further, f_Y evaluates to 1 in both models, which yields the same (uniform) input/output behavior in both models. (To guarantee positivity, we can apply the trick of making a new $f_{Y'}$ such that $f_{Y'}$ returns 0 half the time, and f_Y the other half (i.e., set $f_{Y'} = [f_Y \wedge C]$, where C is a fair coin.) ■

Lemma 9. *There exist values of f_X such that $X, Y P_1(Y|do(X), S = 1) \neq P_2(Y|do(X), S = 1)$.*

Proof. Fix $X = 1, Y = 1$. First notice that f_Z evaluates to U_2 in M_1 and $(U_2 \otimes 1)$ in M_2 . Given that U_2 is uniformly distributed, both quantities coincide (and they represent the effect of X on Z , which is transportable in G). Now the evaluation of f_Y in M_1 reduces to U_1 , while it reduces to 1 in M_2 , which show disagreement and finishes the proof of this Lemma. ■

By Lemma 2, Lemmas 8 and 9 together prove Theorem 8. ■

Remark 3. There exists a different sort of asymmetry in the case of Figure 7(b) (called *sb-graph*), and the nodes X and S do not intersect before meeting Y – i.e., they have disjoint paths and Y lies precisely in their intersection.

Still, this case is not the same of having a *sC-tree* because in *sb-graphs* we need to keep the equality from the S nodes to Y until S intersects X on Y . Employing a similar construct as in the *sp-graph*, we keep the effect of S dormant until it reaches Y and then emerges.

Theorem 9. $P_x^*(y)$ is not transportable in the *sb-graph* (Figure 7(b)).

Proof. We construct two causal models M_1 and M_2 compatible with the *sb-graph* that will agree on $\langle P, P^*, I \rangle$, but disagree on the interventional distribution $P_x^*(y)$.

Let us assume that all variables in $\mathbf{U} \cup \mathbf{V}$ are binary, and let U_1 be the common cause of X and Y , U_2 be the common cause of Z and Y , and U_3 be the random disturbance exclusive to X . Let M_1 and M_2 agree with the following definitions:

$$M_1, M_2 = \begin{cases} X = U_1 \\ Z = ((U_3 \otimes U_2 \otimes 1) \vee S) \otimes (S \wedge U_2) \end{cases}$$

and disagree in respect to Z as follows:

$$\begin{cases} M_1 : Y = Z \otimes U_2 \\ M_2 : Y = X \otimes Z \otimes U_1 \otimes U_2 \end{cases}$$

Both models also agree in respect to $P(\mathbf{U})$, which is defined as follows:

$$P(U_1) = P(U_2) = P(U_3) = 1/2$$

Lemma 10. *The two models agree in the distributions $\langle P, P^*, I \rangle$.*

Proof. Subclaim 1: Let us show that both models agree in the observational and interventional distributions relative to domain Π , i.e., the pair $\langle P, I \rangle$. The index variable S is set to 0 in Π , and $\{X, Z\}$ are defined in the same way in both models, and so it suffices to analyze Y , which in this case evaluates to $(U_3 \otimes 1)$ in both models, preserving the same (uniform) probabilistic behavior. Given that, it is not difficult to see that both models also evaluate in the same way when considering the interventions in I .

Subclaim 2: Let us show that both models agree in the observational distribution P^* relative to Π^* . The index variable S is set 1 in Π^* , given that $\{X, Z\}$ are defined in the same way in both models, together with the uniformity of U make them evaluate in the same way in both models, and Y evaluates to 1 in both models. (As in Lemma 8, the same trick to make the distribution positive could be applied here.) ■

Lemma 11. *There exist values of X, Y such that $P_1(Y|do(X), S = 1) \neq P_2(Y|do(X), S = 1)$.*

Proof. Fix $X = 1, Y = 1$. First notice that f_Z evaluates to $(U_2 \otimes 1)$ in both models, and the evaluation of f_Y in M_1 reduces to 1, while it reduces to U_1 in M_2 . It follows that in M_1 , f_Y evaluates to 1 with probability 1, while in M_2 it evaluates to 1 with probability $P(U_1 = 1)$, which disagree by construction, finishing the proof of this Lemma. ■

By Lemma 2, Lemmas 10 and 11 together prove Theorem 9. ■

Remark 4. There are two complementary components to forge a general scheme to prove arbitrary non-transportability. First, the construct of Theorem 4 shows how to prove non-transportability for general structures such as sC -trees. In the sequel, the specific proofs of non-transportability for the sp -graph (Theorem 9) and sb -graph (Theorem 10) partition the possible interactions between X, S and Y . In the former, X and S intersect before meeting with Y , while in the latter they have disjoint paths and Y lies in their intersection. In the sequel, the proof for the general case combines these analyses, which we show below.

Theorem 4. *Assume there exist F, F' that form a s -hedge for $P_{\mathbf{x}}^*(\mathbf{y})$ in Π and Π^* . Then $P_{\mathbf{x}}^*(\mathbf{y})$ is not transportable from Π to Π^* .*

Proof. We first consider counterexamples with the induced graph $H = De(F)_{G'} \cap An(\mathbf{Y})_{G_x}$, and assume, without loss of generality, that H is a forest. We construct two causal models M_1 and M_2 that will agree on $\langle P, P^*, I \rangle$, but disagree on the interventional distribution $P_{\mathbf{x}}^*(\mathbf{y})$.

Let F be an \mathbf{R} -rooted sC -forest, let \mathbf{V}' be the set of observable variables and \mathbf{U}' be the set of unobservable variables in F . Let us assume that all variables in $\mathbf{U}' \cup \mathbf{V}'$ are binary. Call \mathbf{W} the set of variables pointed by S -nodes in F' , which by the definition of sC -forest is guaranteed to be non-empty.

In model 1, let each $V_i \in \mathbf{V}' \setminus \mathbf{W}$ compute the bit parity of all its observable and unobservable parents (i.e., $f_i^{(1)} = \bigotimes (\bigcup_{V_j \in \mathbf{Pa}_i} V_j)$, where the xor is applied for each element of the set and the result computed so far), while in model 2, let V_i compute the bit parity of all its parents except that any node in F' disregards the parents values if the parent is in F (i.e., $f_i^{(2)} = \bigotimes (\bigcup_{V_j \in \mathbf{Pa}_i \cap F'} V_j)$ if V_i is in F' , and $f_i^{(2)} = f_i^{(1)}$, otherwise).

Define $W \in \mathbf{W}$ as follows:

$$\begin{cases} M_1 : W = ((f_w^{(1)} \otimes U_w^*) \vee S) \otimes (S \wedge (1 \otimes f_w^{(1)})) \\ M_2 : W = ((f_w^{(2)} \otimes U_w^*) \vee S) \otimes (S \wedge (1 \otimes f_w^{(2)})) \end{cases}$$

where f_w is constructed in similar way as f_i in M_1 and M_2 above, and U_w^* is an additional fair coin exclusively pointing to W . Let us call \mathbf{U}_w the collection of such coins. Furthermore, let us assume that each $U_i \in \{\mathbf{U}' \setminus \mathbf{U}_w\}$ is also a fair coin (i.e., $P(U_i) = 1/2$).

Lemma 12. *The two models agree in the distribution of P^* and there exists a value assignment \mathbf{x} for \mathbf{X} such that $P_1(\mathbf{Y} | do(\mathbf{x}), S = 1) \neq P_2(\mathbf{Y} | do(\mathbf{x}), S = 1)$.*

Proof. For $S = 1$, the result follows directly since the systems of equations in both models reduce to the construction given in Theorem 4 at [27]. ■

Lemma 13. *The two models agree in the distributions $\langle P, I \rangle$.*

Proof. Let us show that both models agree in the observational distribution P relative to domain Π . The selection variable S is set to 0 in Π , and note that both systems are the same as in Π^* except that now each variable $W \in \mathbf{W}$ has an extra variable U_w^* pointing to it that should be taken into account in W 's evaluation, and in turn in the whole system.

We have a forest over the endogenous nodes and all functions compute the bit parity of the value of their parents, and so we can view each node as computing the sum mod 2 of its exogenous ancestors in H . We want to show that the distribution of each family is equally likely for each possible assignment (i.e., $P(v_i | \mathbf{pa}_i) = 1/2$, for all v_i, \mathbf{pa}_i).

Let us partition the analysis in two cases. First consider the case of $V_i \in \mathbf{R}$ in which there exists a S -node in the respective sC-tree. Note that the evaluation of V_i relies only on the value of $U_w^* \in \mathbf{U}_w$ in its respective tree since $U \in \{\mathbf{U}' \setminus \mathbf{U}_w\}$ has an even number of endogenous children in F , and it is counted twice, so evaluates to zero (i.e., it does not affect V_i 's evaluation). For now, let us assume that there is only one U_w^* that affects the evaluation of V_i . Given the uniformity of U_w^* , it suffices to show that U_w^* can vary independently for any configuration of the parents of V_i .

For any configuration of $\mathbf{U}' = (U_1 = u_1, \dots, U_w^* = u_w^*, \dots)$, consider the corresponding evaluation of $\mathbf{Pa}_i = \mathbf{pa}_i$, and also $V_i = u_w^*$. We want to show that it is possible to flip the current value of U_w^* from u_w^* to $\neg u_w^*$ while preserving the parents' evaluation \mathbf{pa}_i . Assume this is not so. This implies that the evaluation of \mathbf{Pa}_i and V_i count the same \mathbf{U}' 's, contradiction.

To see why, consider $\mathbf{Pa}_i^* \subseteq \mathbf{Pa}_i$ the set of parents of V_i that are descendents of U_w^* . Now, for each of these parents flip the minimum number of variables from $\mathbf{U} \setminus \mathbf{U}_w$, and call this set \mathbf{U}^* . (Note that this is always possible since we need at most one U for each parent, which should exist by construction of sC-forest.) Now, make $U_w^* = \neg u_w^*$, and note that $\mathbf{Pa}_i = \mathbf{pa}_i$ since flipping the values of \mathbf{U}^* compensates the flip of U_w^* . But it is also true now that V_i evaluates to $\neg u_w^*$ since, in the same way as before, all other variables in $\{\mathbf{U} \setminus \mathbf{U}_w\}$ are cancelled out in V_i 's evaluation, including the ones in \mathbf{U}^* . This proves the claim.

Consider the following two facts: **Subclaim 1:** Let X and Y be two binary variables such that $P(X = x) = p \neq 1/2$ and $P(Y = y) = q = 1/2$. Then the probabilistic input/output behavior of $Z = \text{XOR}(X, Y)$ is the same of Y . The variable $Z = 1$ whenever $\{(X = 1, Y = 0), (X = 0, Y = 1)\}$, which happens with probability $pq + (1 - p)(1 - q)$. Since $q = 1/2$, the expression reduces to $p * 1/2 + (1 - p) * 1/2 = 1/2$.

Subclaim 2: Let X and Y be two binary variables such that $P(X = x) = P(Y = y) = p = 1/2$. Then the probabilistic input/output behavior of $Z = XOR(X, Y)$ is the same of X (or Y). This follows directly from Subclaim 1. It is clear that if there are multiple nodes from \mathbf{U}_w in the evaluation of V_i , the same construction is also valid given the subclaim above. It is also not difficult to generalize this argument to consider root set that are not singleton, including roots in which there are not S -nodes as ancestors.

Finally, let us consider the case of $V_i \in \{F \setminus \mathbf{R}\}$. It suffices to show that the function from $\mathbf{U} \setminus \mathbf{U}_w$ to $\mathbf{V} \setminus \mathbf{R}$ is 1–1 when we fix $\mathbf{U}_w = \mathbf{u}_w$. We use the same argument as Shpitser. Assume this is not so, and fix two instantiations of $\mathbf{U} \setminus \mathbf{U}_w$ that map to the same value of $\mathbf{V} \setminus \mathbf{R}$, and differ by the set $\mathbf{U}^* = \{U_1, \dots, U_k\}$. Since the bidirected edges form a spanning tree, there exists \mathbf{V}^* with an odd number of parents in \mathbf{U}^* (and were not in \mathbf{R} , by construction). Order them topologically and let the topmost be called X . Note that if we flip all values in \mathbf{U}^* , the value of X will also flip, contradiction. Given the uniformity of \mathbf{U}' , the claim follows. We can put this together with the previous claim, and the result follows. We can add fair coins as the input to all other variables outside F , which will imply the claim for the whole graph G .

In regard to the equality between I , note that given that the equality of both models holds for P , and removing edges due to interventions will just make some nodes from $\mathbf{U} \setminus \mathbf{U}_w$ to have an odd number of children, it is not difficult to see based on the previous argument that this just creates more variables that are free to vary, which will entail the same probabilistic uniform behavior in both models. Another way to see this fact is to consider the new exogenous variables from $\{\mathbf{U} \setminus \mathbf{U}_w\}$ that have only one children after the intervention as analogous to U_w^* , and so the same argument follows. ■

Finally, Lemma 2 together with Lemmas 12 and 13 prove Theorem 4. ■

Theorem 5 (soundness). *Whenever **sID** returns an expression for $P_x^*(\mathbf{y})$, it is correct.*

Proof. Noting that the selection diagram inputted to **sID** is also a causal diagram over Π^* , and trivial transportability is equivalent to identifiability in Π^* , the correctness of the identifiability calls was already established elsewhere [27, 34].

It remains to show the correctness of the test in line 10 of **sID**. First note that, by construction, \mathbf{X}' in each local call is always a set of pre-treatment covariates. But now the correctness follows directly by S -admissibility of \mathbf{X}' together with Corollary 1 in Ref. 1. Further note that the set of \mathbf{Z} -nodes outside the local component will not affect separability of the S -nodes inside it (following the topology of the hedge), and other S -nodes outside can be removed from the expression before the test. More specifically, note that the effect Q^* in each local call that uses line 10 can be expressed in its expanded form (using a typical C -component decomposition), and given that the independence imposed by S -admissibility holds, together with the fact that both populations share the same causal graph G , allow that the functions of Π^* to be replaced with the respective functions in Π , which implies the result. ■

Remark 5. The next results are similar to the identification counterparts given in Refs. 26, 69.

Theorem 6. *Assume **sID** fails to transport $P_x^*(\mathbf{y})$ (executes line 11). Then there exists $\mathbf{X}' \subseteq \mathbf{X}$, $\mathbf{Y}' \subseteq \mathbf{Y}$, such that the graph pair D, C_0 returned by the fail condition of **sID** contain as edge subgraphs sC -forests F, F' that form a s -hedge for $P_x^*(\mathbf{y}')$.*

Proof. Before failure **sID** evaluated false consecutively at lines 5, 6, and 10, so D local to this call is a sC -component, and let \mathbf{R} be its root set. We can remove some directed arrows from D while preserving \mathbf{R} as root, yielding a \mathbf{R} -rooted sC -forests F . Since by construction $F' = F \cap C_0$ is closed under descendants and only directed arrows were removed, both F, F' are sC -forests. Also by construction, $\mathbf{R} \subset An(\mathbf{Y})_{D_x}$ together with the fact that \mathbf{X} and \mathbf{Y} from the recursive call are clearly subsets of the original input, finish the proof.

Corollary 3 (completeness). ***sID** is complete.*

Proof. The result follows from Theorem 6 where $P_{\mathbf{x}'}^*(\mathbf{y}')$ is not transportable in H . But now, it is easy to add the remaining variables from G , making them independent of H (e.g., as random coins). So, the models in the counterexample induce G , and witness the non-transportability of $P_{\mathbf{x}}^*(\mathbf{y})$.

Corollary 4. $P_{\mathbf{x}}^*(\mathbf{y})$ is transportable from Π to Π^* in G if and only if there is not s -hedge for $P_{\mathbf{x}'}^*(\mathbf{y}')$ in G for any $\mathbf{X}' \subseteq \mathbf{X}$ and $\mathbf{Y}' \subseteq \mathbf{Y}$.

Proof. Follows directly from the previous Corollary. ■

Theorem 7. The rules of *do*-calculus, together with standard probability manipulations are complete for establishing transportability of all effects of the form $P_{\mathbf{x}}^*(y)$.

Proof. It was shown elsewhere [69] that the steps of **sID** but line 10 correspond to sequences of standard probability manipulations and applications of the rules of *do*-calculus. The line 10 is constituted by a conditional independence judgment, and standard probability operations for the replacement of the functions based on the invariance allowed by the S -admissibility of the local X' in each recursive call (as discussed above in the proof of correctness). ■

References

1. Pearl J, Bareinboim E. Transportability of causal and statistical relations: a formal approach. In Proceedings of the Twenty-Fifth National Conference on Artificial Intelligence (AAAI 2011). Menlo Park, CA: AAAI Press, 2011:247–54.
2. Bareinboim E, Pearl J. Causal transportability with limited experiments. In Proceedings of the Twenty-Seventh AAAI Conference on Artificial Intelligence (AAAI 2013), Menlo Park, CA: AAAI Press, 2013, forthcoming.
3. Bareinboim E, Pearl J. Meta-transportability of causal effects: A formal approach. In Proceedings of the Sixteenth International Conference on Artificial Intelligence and Statistics (AISTATS 2013), 2013, forthcoming.
4. Campbell D, Stanley J. Experimental and quasi-experimental designs for research. Chicago: Wadsworth Publishing, 1963.
5. Manski C. Identification for prediction and decision. Cambridge, Massachusetts: Harvard University Press, 2007.
6. Glass GV. Primary, secondary, and meta-analysis of research. *Educ Res* 1976;5:3–8.
7. Hedges LV, Olkin I. Statistical methods for meta-analysis. Orlando, FL: Academic Press, 1985.
8. Owen AB. Karl pearsons meta-analysis revisited. *Ann Stat* 2009;37:3867–92.
9. Höfler M, Gloster A, Hoyer J. Causal effects in psychotherapy: counterfactuals counteract overgeneralization. *Psychother Res* 2010, DOI: 10.1080/10503307.2010.501041.
10. Shadish W, Cook T, Campbell D. Experimental and Quasi-Experimental Designs for Generalized Causal Inference. Boston: Houghton-Mifflin, 2nd ed., 2002.
11. Adelman L. Experiments, quasi-experiments, and case studies: a review of empirical methods for evaluating decision support systems. *Systems, Man and Cybernetics, IEEE Transactions on*, 1991;21:93–301.
12. Morgan S, Winship C. Counterfactuals and causal inference: methods and principles for social research (Analytical Methods for Social Research). New York: Cambridge University Press, 2007.
13. Pearl J. Causal diagrams for empirical research. *Biometrika* 1995;82:669–710.
14. Greenland S, Pearl J, Robins J. Causal diagrams for epidemiologic research. *Epidemiology* 1999;10:37–48.
15. Spirtes P, Glymour C, Scheines R. Causation, prediction, and search. Cambridge, MA: MIT Press, 2nd ed., 2001.
16. Pearl J. Causality: models, reasoning, and inference. New York: Cambridge University Press, 2nd ed., 2009.
17. Koller D, Friedman N. Probabilistic graphical models: principles and techniques. Cambridge, MA: MIT Press, 2009.
18. Westergaard H. Scope and method of statistics. *Am Stat Assoc* 1916;15:229–76.
19. Yule G. On some points relating to vital statistics, more especially statistics of occupational mortality. *J R Stat Soc* 1934;97:1–84.
20. Lane P, Nelder J. Analysis of covariance and standardization as instances of prediction. *Biometrics* 1982;38:613–21.
21. Cole S, Stuart E. Generalizing evidence from randomized clinical trials to target populations. *Am J Epidemiol* 2010;172: 107–15.
22. Pearl J. Causality: models, reasoning, and inference. New York: Cambridge University Press, 2nd ed., 2000.
23. Pearl J. Causal inference in statistics: an overview. *Stat Surv* 2009;3:96–146.
24. Pearl J, Verma T. A theory of inferred causation. In Allen J, Fikes R, Sandewall E, editors. Principles of knowledge representation and reasoning: Proceedings of the Second International Conference. San Mateo, CA: Morgan Kaufmann, 1991:441–52.

25. Bareinboim E, Brito C, Pearl J. Local characterizations of causal bayesian networks. In Croitoru M, Corby O, Howse J, Rudolph S, Wilson N, editors. GKR-IJCAI, Lecture Notes in Artificial Intelligence (7205), Springer-Verlag, 2012:1–17.
26. Tian J, Pearl J. A general identification condition for causal effects. In Proceedings of the Eighteenth National Conference on Artificial Intelligence (AAAI 2002). Menlo Park, CA: AAAI Press/The MIT Press, 2002:567–73.
27. Shpitser I, Pearl J. Identification of joint interventional distributions in recursive semi-Markovian causal models. In Proceedings of the Twenty-First National Conference on Artificial Intelligence (AAAI 2006). Menlo Park, CA: AAAI Press, 2006:1219–26.
28. Spirtes P, Glymour C, Scheines R. Causation, prediction, and search. New York: Springer-Verlag, 1993.
29. Galles D, Pearl J. Testing identifiability of causal effects. In Besnard P, Hanks S, editors. Proceedings of the Eleventh Conference on Uncertainty in Artificial Intelligence (UAI 1995). San Francisco: Morgan Kaufmann, 1995:185–95.
30. Pearl J, Robins J. Probabilistic evaluation of sequential plans from causal models with hidden variables. In Besnard P, Hanks S, editors. Proceedings of the Eleventh Conference on Uncertainty in Artificial Intelligence (UAI 1995). San Francisco: Morgan Kaufmann, 1995:444–53.
31. Halpern J. Axiomatizing causal reasoning. In Cooper G, Moral S, editors. Uncertainty in artificial intelligence. San Francisco, CA: Morgan Kaufmann, 1998:202–10, also *J Artif Intell Res* 2000;12:3, 17–37.
32. Kuroki M, Miyakawa M. Identifiability criteria for causal effects of joint interventions. *J R Stat Soc* 1999;29:105–17.
33. Verma T, Pearl J. Equivalence and synthesis of causal models. In Proceedings of the Sixth Conference on Uncertainty in Artificial Intelligence (UAI 1990). Cambridge, MA, 1990:220–27, also in Bonissone P, Henrion M, Kanal LN, Lemmer JF, editors. Uncertainty in artificial intelligence 6. Amsterdam, The Netherlands: Elsevier Science Publishers, B.V., 1990:255–68, 1991.
34. Huang Y, Valtorta M. Identifiability in causal bayesian networks: A sound and complete algorithm. In Proceedings of the Twenty-First National Conference on Artificial Intelligence (AAAI 2006). Menlo Park, CA: AAAI Press, 2006:1149–56.
35. Pearl J. Direct and indirect effects. In Proceedings of the Seventeenth Conference on Uncertainty in Artificial Intelligence (UAI 2001). San Francisco, CA: Morgan Kaufmann, 2001:411–20.
36. Pearl J. The mediation formula: A guide to the assessment of causal pathways in nonlinear models. In Berzuini C, Dawid P, Bernardinell L, editors. Causality: statistical perspectives and applications. New York: Wiley, Chapter 12, 2012.
37. Bareinboim E, Pearl J. Causal inference by surrogate experiments: z-identifiability. In de Freitas N, Murphy K, editors. Proceedings of the Twenty-Eighth Conference on Uncertainty in Artificial Intelligence (UAI 2012), AUAI Press, 2012, 113–20.
38. Cornfield J. A method of estimating comparative rates from clinical data; applications to cancer of the lung, breast, and cervix. *J Natl Cancer Inst* 1951;11:1269–75.
39. Whittemore A. Collapsibility of multidimensional contingency tables. *J R Stat Soc, B* 1978;40:328–40.
40. Geng Z, Guo J, Fung W-K. Criteria for confounders in epidemiological studies. *J Royal Stat Soc Series B* 2002;64:3–15.
41. Heckman JJ. Sample selection bias as a specification error. *Econometrica* 1979;47:153–61.
42. Robins JM, Hernan M, Brumback B. Marginal structural models and causal inference in epidemiology. *Epidemiology* 2000;11:550–60.
43. Hernán M, Hernández-Díaz S, Robins J. A structural approach to selection bias. *Epidemiology* 2004;15:615–25.
44. Lauritzen SL, Richardson TS. Discussion of mccullagh: sampling bias and logistic models. *J R Stat Soc Ser B* 2008;70:140–50.
45. Geneletti S, Richardson S, Best N. Adjusting for selection bias in retrospective, case-control studies. *Biostatistics* 2009;10:17–31.
46. Weisberg H, Hayden V, Pontes V. Selection criteria and generalizability within the counterfactual framework: explaining the paradox of antidepressant-induced suicidality? *Clin Trials* 2009;6:109–18.
47. Stuart EA, Cole SR, Bradshaw CP, Leaf PJ. The use of propensity scores to assess the generalizability of results from randomized trials. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 2011;174:369–86.
48. Angrist J, Imbens G, Rubin D. Identification of causal effects using instrumental variables (with comments). *J Am Stat Assoc* 1996;91:444–72.
49. Didelez V, Kreiner S, Keiding N. Graphical models for inference under outcome-dependent sampling. *Stat Sci* 2010;25:368–87.
50. Bareinboim E, Pearl J. Controlling selection bias in causal inference. In Girolami M, Lawrence N, editors. Proceedings of The Fifteenth International Conference on Artificial Intelligence and Statistics (AISTATS 2012), *JMLR* (22), 2012:100–08.
51. Pearl J. A solution to a class of selection-bias problems. Technical Report R-405, Cognitive Systems Laboratory, Department of Computer Science, UCLA, 2012.
52. Hernán M, VanderWeele T. Compound treatments and transportability of causal inference. *Epidemiology* 2011;22:368–77.
53. Petersen M. Compound treatments, transportability, and the structural causal model: the power and simplicity of causal graphs. *Epidemiology* 2011;22:378–81.
54. Cox D. The Planning of Experiments. New York: John Wiley and Sons, 1958.
55. Heckman J. Randomization and social policy evaluation. In Manski C, Garfinkle I, editors. Evaluations: welfare and training programs. Cambridge, MA: Harvard University Press, 1992:201–30.
56. Hotz VJ, Imbens G, Mortimer JH. Predicting the efficacy of future training programs using past experiences at other locations. *J Econ* 2005; 125:241–70.

57. Pearl J, Bareinboim E. Transportability of causal and statistical relations: A formal approach. Technical Report Technical Report r-372, Cognitive Systems Laboratory, Department of Computer Science, UCLA, 2011.
58. Pearl J. Some thoughts concerning transfer learning with applications to meta-analysis and data sharing estimation. Technical Report R-387, Cognitive Systems Laboratory, Department of Computer Science, UCLA, 2012.
59. Bareinboim E, Pearl J. Transportability of causal effects: completeness results. In Hoffmann J, Selman B, editors. Proceedings of The Twenty-Sixth Conference on Artificial Intelligence (AAAI 2012), 2012:698–704.
60. Bollen KA, Pearl J. Eight myths about causality and structural equation models. In Morgan SL, editor. Handbook of Causal Analysis for Social Research (in press), New York: Springer, 2013, Chapter 15.
61. Haavelmo T. The statistical implications of a system of simultaneous equations. *Econometrica* 1943;11:1–12, reprinted in D. F. Hendry DF, Morgan MS, editors. The foundations of econometric analysis. Cambridge University Press, 1995:477–90.
62. Strotz R, Wold H. Recursive versus nonrecursive systems: an attempt at synthesis. *Econometrica* 1960;28:417–27.
63. Pearl J. Trygve Haavelmo and the emergence of causal calculus. Technical Report R-391, Cognitive Systems Lab, Department of Computer Science, UCLA; To appear: *Econometric Theory*, special issue on Haavelmo Centennial, 2012.
64. Lehmann EL, Casella G. Theory of point estimation (Springer Texts in Statistics). New York: Springer, 2nd ed., 1998.
65. Pearl J. Probabilistic reasoning in intelligent systems. San Mateo, CA: Morgan Kaufmann, 1988.
66. Hayduk L, Cummings G, Stratkotter R, Nimmo M, Grygoryev K, Dosman D, et al. Pearls d-separation: one more step into causal thinking. *Struct Equ Modeling* 2003;10:289–311.
67. Glymour M, Greenland S. Causal diagrams. In Rothman K, Greenland S, Lash T, editors. *Modern epidemiology*. Philadelphia, PA: Lippincott Williams & Wilkins, 3rd ed., 2008:183–209.
68. Berkson J. Limitations of the application of fourfold table analysis to hospital data. *Biometrics Bull* 1946;2:47–53.
69. Shpitser I, Pearl J. Identification of conditional interventional distributions. In Dechter R, Richardson T, editors. Proceedings of the Twenty-Second Conference on Uncertainty in Artificial Intelligence (UAI 2006). Corvallis, OR: AUAI Press, 2006:437–44.
70. Huang Y, Valtorta M. Pearl's calculus of intervention is complete. In Dechter R, Richardson T, editors. Proceedings of the Twenty-Second Conference on Uncertainty in Artificial Intelligence. Corvallis, OR: AUAI Press; 2006:217–24.
71. Tian J. Studies in causal reasoning and learning. PhD Thesis, Computer Science Department, University of California, Los Angeles, CA, 2002.

